

ペンタクォーク探索実験 J-PARC E19: 2nd Run Result

京大理、他
森津 学

for the J-PARC E19 collaboration



Contents

- 1. Introduction**
- 2. J-PARC E19 experiment**
 - (1st run result)**
- 3. E19-2nd run**
 - analysis status and preliminary result**
- 4. Summary**


Pentaquark Θ^+

- Five-quark state (qqqq \bar{q})
 - Allowed combination by QCD.
- Θ^+ includes \bar{s} . \rightarrow At least 5-quark components.
- Θ^+ has extraordinary narrow width
 - despite of its mass(~ 1530) over the KN threshold. \rightarrow Need some mechanism to suppress decay.
 - Diquark-triquark configuration ?
 - Diquark correlation ?
 -

✓ Pentaquark Θ^+ must offer us a good opportunity to understand low energy QCD dynamics (, if exist) .



Karliner, Lipkin,
PLB 575, 249 (2003).



Jaffe, Wilczek,
PRL 91, 232003 (2003).
3

A Lot of Θ^+ Searches

Group	Reaction	Mass (MeV)	Width (MeV)	Statistical significance (σ)
LEPS	$\gamma C \rightarrow K^+ K^- (n)$	1540 ± 10	< 25	4.6
LEPS	$\gamma C \rightarrow K^+ K^- (n)$	1524 ± 2	< 25	5.1
DIANA	$K^+ X e \rightarrow K_s^0 p X$	1539 ± 2	< 9	4.4
DIANA	$K^+ X e \rightarrow K_s^0 p X$	1538 ± 2	0.39 ± 0.1	8
CLAS(d)	$\gamma d \rightarrow K^+ K^- p (n)$	1542 ± 5	< 21	(5.2)
CLAS(p)	$\gamma p \rightarrow \pi^+ K^+ K^- (n)$	1555 ± 10	< 26	7.8
SAPHIR	$\gamma p \rightarrow K^+ K_s^0 n X$	1540 ± 6	< 25	4.8
ITEP	$\nu A \rightarrow K_s^0 p X$	1533 ± 5	< 20	6.7
HERMES	$e^+ d \rightarrow K_s^0 p X$	1528 ± 3	12 ± 9	4.2
COSY-TOF	$pp \rightarrow K_s^0 p \Sigma^+$	1530 ± 5	< 18	4.7
ZEUS	$e^+ p \rightarrow e^+ K_s^0 p X$	1522 ± 3	8 ± 4	4.6
NOMAD	$\nu A \rightarrow K_s^0 p X$	1529 ± 3	$2 \sim 3$	4.3
SVD	$p A \rightarrow K_s^0 p X$	1526 ± 5	< 24	5.6
SVD	$p A \rightarrow K_s^0 p X$	1523 ± 5	< 14	8.0

Positive results

Negative results

Group	Reaction	Limit
BES	$e^+ e^- \rightarrow J/\Psi \rightarrow \Theta \bar{\Theta}$	$< 1.1 \times 10^{-5}$ B.R. (90% C.L.)
BES	$e^+ e^- \rightarrow \Psi(2S) \rightarrow \Theta \bar{\Theta}$	$< 8.4 \times 10^{-6}$ B.R. (90% C.L.)
ALEPH	$e^+ e^- \rightarrow Z \rightarrow p K_s^0 X$	$< 6.2 \times 10^{-4}$ B.R. (95% C.L.)
BarBar	$e^+ e^- \rightarrow \Upsilon(4S) \rightarrow p K_s^0 X$	$< 1.0 \times 10^{-4}$ B.R. (90% C.L.)
BarBar	$e B e \rightarrow p K_s^0 X$	not given
Belle	$e^+ e^- \rightarrow B^0 \bar{B}^0 \rightarrow p \bar{p} K_s^0 X$	$< 2.3 \times 10^{-7}$ B.R. (90% C.L.)
Belle	$K^+ n \rightarrow K_s^0 p X$	$\Gamma < 0.64 \text{ MeV}$ (90% C.L.)
CDF	$p \bar{p} \rightarrow K_s^0 p X$	$< 0.03 \times \Lambda^*$ (90% C.L.)
SPHINX	$p C \rightarrow K_s^0 p X$	$< 0.1 \times \Lambda^*$ (90% C.L.)
HERA-B	$p A \rightarrow K_s^0 p X$	$< 2.7\% \times \Lambda^*$ (95% C.L.)
HyperCP	$p C u \rightarrow K_s^0 p X$	$< 0.3\% K_s^0 p$
FOCUS	$\gamma B e O \rightarrow K_s^0 p X$	$< 0.02 \times \Sigma^*$ (95% C.L.)
PHENIX	$d A u \rightarrow K^- \bar{n} X$	not given
WA89	$\Sigma^+ A \rightarrow K_s^0 p X$	$< 1.8 \mu\text{b/A}$ (99% C.L.)
CLAS	$\gamma p \rightarrow \bar{K}_s^0 K^+ n$	$< 0.8 \text{ nb}$ (95% C.L.)
CLAS	$\gamma d \rightarrow K^- p K^+ n$	$< 0.15 - 3 \text{ nb}$ (95% C.L.)
CLAS	$\gamma d \rightarrow K^+ n \Lambda$	$< 5 - 25 \text{ nb}$ (95% C.L.)
COSY-TOF	$pp \rightarrow \Sigma^+ p K_s^0$	$< 0.15 \mu\text{b/A}$ (95% C.L.)
NOMAD	$\nu A \rightarrow K_s^0 p X$	$< 2.13 \times 10^{-3} \nu\text{CC}$ (90% C.L.)

A Lot of Θ^+ Searches

Group	Reaction	Mass (MeV)	Width (MeV)	Statistical significance (σ)
LEPS	$\gamma C \rightarrow K^+ K^- (n)$	1540 ± 10	< 25	4.6
LEPS	$\gamma C \rightarrow K^+ K^- (n)$	1524 ± 2	< 25	5.1
DIANA	$K^+ X e$			
DIANA	$K^+ X e$			
CLAS(d)	$\gamma d \rightarrow K^+ n$			
CLAS(p)	$\gamma p \rightarrow \pi^+ n$			
SAPHIR	$\gamma p \rightarrow K^+ n$			
ITEP	$\nu A \rightarrow \Lambda^+ K^-$			
HERMES	$e^+ d \rightarrow e^+ \Lambda^+ K^-$			
COSY-TOF	$pp \rightarrow K_s^0 p \Lambda^+$	1538 ± 5	< 15	4.7
ZEUS	$e^+ p \rightarrow e^+ K_s^0 p X$	1522 ± 3	8 ± 4	4.6
NOMAD	$\nu A \rightarrow K_s^0 p X$	1529 ± 3	$2 \sim 3$	4.3
SVD	$p A \rightarrow K_s^0 p X$	1526 ± 5	< 24	5.6
SVD	$p A \rightarrow K_s^0 p X$	1523 ± 5	< 14	8.0

✓ *Not well established in experiments*
 → *“Must confirm the existence/non-existence of Θ^+ at first”*

ve results

Group	Reaction	Limit
BES	$e^+ e^- \rightarrow J/\Psi \rightarrow \Theta \bar{\Theta}$	$< 1.1 \times 10^{-5}$ B.R. (90% C.L.)
BES	$e^+ e^- \rightarrow \Psi(2S) \rightarrow \Theta \bar{\Theta}$	$< 8.4 \times 10^{-6}$ B.R. (90% C.L.)
ALEPH	$e^+ e^- \rightarrow Z \rightarrow p K_s^0 X$	$< 6.2 \times 10^{-4}$ B.R. (95% C.L.)
BarBar	$e^+ e^- \rightarrow \Upsilon(4S) \rightarrow p K_s^0 X$	$< 1.0 \times 10^{-4}$ B.R. (90% C.L.)

Positive results



✓ *Low energy hadronic reaction (π or K beam)*


- Few data
- Expect sizable production cross section.
- Complementary to the photo-production.

WA89	$\Sigma^+ A \rightarrow K_s^+ p \Lambda$	$< 1.8 \mu\text{b}/A$ (99% C.L.)
CLAS	$\gamma p \rightarrow \bar{K}_s^0 K^+ n$	< 0.8 nb (95% C.L.)
CLAS	$\gamma d \rightarrow K^- p K^+ n$	$< 0.15 - 3$ nb (95% C.L.)
CLAS	$\gamma d \rightarrow K^+ n \Lambda$	$< 5 - 25$ nb (95% C.L.)
COSY-TOF	$pp \rightarrow \Sigma^+ p K_s^0$	$< 0.15 \mu\text{b}/A$ (95% C.L.)
NOMAD	$\nu A \rightarrow K_s^0 p X$	$< 2.13 \times 10^{-3} \nu\text{CC}$ (90% C.L.)

Θ^+ search by high-resolution spectroscopy via $\pi^- + p \rightarrow \Theta^+ + K^-$: J-PARC E19

Previous KEK-PS E522 experiment

- Is this a sign of Θ^+ ?
- Not enough sensitivity
- They did not conclude the evidence of Θ^+ .
- mass resolution
 $\Delta M \sim 13.4 \text{ MeV (FWHM)}$



J-PARC E19 experiment

- same reaction as E522
- High resolution : SKS → $\Delta M < 2 \text{ MeV (FWHM)}$
- High statistics : High intensity beam at J-PARC

⇒ **Conclusive result by higher sensitivity.**

The first physics run at the J-PARC hadron facility !

Experimental setup

K1.8 beam line spectrometer & SKS

⇒ Missing mass spectroscopy

➤ K1.8 beam line spectrometer : p_π

PID counters

- Timing counters : TOF
- Gas Cherenkov (π/e) : $n=1.002$

Tracking

- MWPCs : 1 mm pitch
- MWDCs : 3 mm pitch

➤ SKS system : p_K

PID counters


- Timing counter
- Aerogel Cherenkov (K/π) : $n=1.05$
- Lucite Cherenkov (K/p) : $n=1.49$

Tracking

- MWDCs : 3 mm pitch
- DCs : 10 mm pitch, $2\text{ m} \times 1\text{ m}$ size

➤ Target: Liquid hydrogen

- $\sim 0.86\text{ g/cm}^2$
- Free from Fermi motion effect



History of E19

0. 2009/10 ~

- K1.8 beam line & detector commissioning start.

1. 1st run (2010/10-11)

- examine the 2.6σ bump structure observed in E522 at $p_\pi = 1.92 \text{ GeV/c}$.
- accumulated 7.8×10^{10} of beam π on target.

□ Earthquake (2011/3)

- Realignment of all the detectors and magnets including SKS.

2. 2nd run (2012/2)


- new data at the highest beam momentum of 2 GeV/c .
- accumulated 8.7×10^{10} of beam π on target.

Successful completion of both 1st and 2nd run


1st run result of E19

accepted in PRL,
arxiv.1203.3604 [nucl-ex]

$\pi^- + p \rightarrow K^- + X$ @ 1.92 GeV/c



- No prominent peak structure
- Upper limit: $< 0.26 \mu\text{b/sr}$
@ 1.51–1.55 GeV/c²



- ✓ s-channel dominance
- ✓ $\Gamma_{\Theta} \propto g^2_{KN\Theta} \propto \sigma_{\text{tot}}$
→ Upper limit of decay width


- 0.72 MeV for $\frac{1}{2}^+$
- 3.1 MeV for $\frac{1}{2}^-$

2nd run of E19

Theoretical calculations :

Hyodo, Hosaka, PRC 72, 055202 (2005).

- Beam time: 2012/Feb
 - Higher beam momentum
2.0 GeV/c (= Max. of K1.8 B.L.)
 - Expecting increased cross section
→ **higher sensitivity**
- **Stringent restriction
on the Θ^+ production
via these hadronic reactions.**



Rotation of SKS




Injection angle : 30 \rightarrow 15 deg.


- *Higher-momentum acceptance.*
- *For future experiments.*

Setup difference of 1st and 2nd run

E19-1st run (2010)



E19-2nd run (2012)



K1.8 Beam spectrometer

Beam Momentum

➤ K1.8 beam line spectrometer : p_π


PID counters

- Timing counters : TOF
- Gas Cherenkov (π/e) : $n=1.002$


Tracking

- MWPCs : 1 mm pitch
- MWDCs : 3 mm pitch


Beam mom. of
2 GeV/c is well
reconstructed.



p_{Beam} [GeV/c]



Beam TOF



π is clearly identified
using TOF btw 2 sets
of hodoscopes

Beam TOF [ns] 13

SKS spectrometer


➤ SKS system : p_K

PID counters

- Timing counter
- Aerogel Cherenkov (K/ π) : $n=1.05$
- Lucite Cherenkov (K/p) : $n=1.49$


Tracking

- MWDCs : 3 mm pitch
- DCs : 10 mm pitch, 2m \times 1m size




We can
separate only K
very clearly.

Scattered particle M^2




**Good momentum
reconstruction and PID !!**

Vertex Reconstruction




$\phi 67.8 \times 120 \text{ mm}$

Vertex-(X vs Y)



Consistent with horizontally oblate beam shape.

Vertex-Z



Target cell is clearly identified !!

Performance of the spectrometers

Calibration

- $\pi^+ + p \rightarrow K^+ + \Sigma^+$ @ 1.37 GeV/c
- Missing mass resolution:
 $\Delta M_{\Sigma} = 2.0 \text{ MeV (FWHM)}$
Equivalent to the 1st run !!
Cf.) $\Delta M_{\Sigma} = 1.9 \pm 0.1 \text{ MeV @ E19-1st}$


⇒ estimate Θ^+ case:

$\Delta M_{\Theta} = 1.75 \text{ MeV (FWHM)}$

- Yield estimation (rough):

Almost Consistent with the 1st run !!

Σ^+ Missing Mass




$$\Gamma = 2.02 \pm 0.06$$

Enough performance !!

Preliminary result of E19-2nd run

Missing Mass : $p(\pi^-, K^-)X$ @ $p_{\pi} = 2.0$ GeV/c



- Analysis parameters were not finally tuned yet.
- No clear peak structure was observed.
- Evaluation of efficiency is on-going.
- Tentative expected sensitivity $\sim 0.3 \mu\text{b/sr}$.

Summary

- **J-PARC E19 : High-resolution search** via $\pi^- p \rightarrow K^- \Theta^+$ reaction
 - The first physics experiment at the J-PARC hadron facility !
 - 1st run result was accepted in PRL. (@ 1.92 GeV/c beam)
 - More than 10 times higher sensitivity than E522.
 - No clear Θ^+ peak $\rightarrow < 0.26 \mu\text{b/sr}$
 - Strong constraint : $\Gamma < \sim 1 \text{ MeV}$
- 2nd run was successfully carried out. (@ 2 GeV/c beam)
 - Good performance of both K1.8BS and SKS.
 - No clear Θ^+ peak (preliminary)
 - Efficiency evaluation etc. are on-going.